

AUGUST 2021

EXECUTIVE SUMMARY

SISKIYOU COUNTY FLOOD CONTROL & WATER
CONSERVATION DISTRICT

Shasta Valley Groundwater Sustainability Plan

PUBLIC DRAFT REPORT



**SISKIYOU COUNTY FLOOD CONTROL AND WATER CONSERVATION DISTRICT
GROUNDWATER SUSTAINABILITY AGENCY
SHASTA VALLEY GROUNDWATER SUSTAINABILITY PLAN**

BOARD

Brandon Criss, County of Siskiyou
Ed Valenzuela, County of Siskiyou
Michael Kobseff, County of Siskiyou
Nancy Ogren, County of Siskiyou
Ray A. Haupt, County of Siskiyou

STAFF

Matt Parker, Natural Resources Specialist, County of Siskiyou

TECHNICAL TEAM

Laura Foglia, LWA
Thomas Harter, UC Davis

Andrew Calderwood, UC Davis
Brad Gooch, UC Davis
Cab Esposito, LWA
Katrina Arredondo, LWA
Kelsey McNeill, LWA
Claire Kouba, UC Davis
Bill Rice, UC Davis

ADVISORY COMMITTEE

John Tannaci, Chair, Residential
Blair Hart, Private Pumper
Gregg Werner, Environmental/Conservation
Justin Holmes, Edson Foulke Ditch Company
Pete Scala, Private Pumper
Grant Johnson, Tribal Representative
Tristan Allen, Montague Water Conservation District
Steve Mains, Grenada Irrigation District
Robert Moser, Municipal/City
Lisa Faris, Big Springs Irrigation District
Justin Sandahl, Shasta River Water Users Association

1 **Executive Summary**

2 **ES-1: INTRODUCTION (CHAPTER 1)**

3 **Background (Section 1.1)**

4 *Section 1 describes the Sustainable Groundwater Management Act and the purpose of*
5 *the Groundwater Sustainability Plan. Section 1 also introduces the management structure*
6 *of the agencies developing and implementing the GSP.*

7 The 2014 Sustainable Groundwater Management Act (SGMA) was established to provide
8 local and regional agencies the authority to sustainably manage groundwater resources
9 through the development and implementation of GSPs for high and medium priority
10 subbasins (e.g., Shasta Valley). In accordance with SGMA, this GSP was developed and
11 will be implemented by the Siskiyou County Flood Control and Water Conservation
12 District, the GSA representing the Basin.

13 The California Department of Water Resources (DWR) and the State Water Resources
14 Control Board (State Board) provide primary oversight for implementation of SGMA. DWR
15 adopted regulations that specify the components and evaluation criteria for groundwater
16 sustainability plans, alternatives to Groundwater Sustainability Plans (GSPs), and
17 coordination agreements to implement such plans. To satisfy the requirements of SGMA,
18 local agencies must do the following:

19 Locally controlled and governed Groundwater Sustainability Agencies (GSAs) must be
20 formed for all high- and medium-priority groundwater basins in California.

- 21 • GSAs must develop and implement GSPs or Alternatives to GSPs that define a
22 roadmap for how groundwater basins will reach long-term sustainability.
- 23 • The GSPs must consider six sustainability indicators defined as: groundwater level
24 decline, groundwater storage reduction, seawater intrusion, water quality
25 degradation, land subsidence, and surface-water depletion.
- 26 • GSAs must submit annual reports to DWR each April 1 following adoption of a
27 GSP.
- 28 • Groundwater basins should reach sustainability within 20 years of implementing
29 their GSPs.

30 This GSP was prepared to meet the regulatory requirements established by DWR, as
31 shown in the completed GSP Elements Guide, provided in Appendix 1-D, which is
32 organized according to the California Code of Regulation Sections of the GSP Emergency
33 Regulations.

34 **Purpose of the Groundwater Sustainability Plan**

35 The Shasta Valley GSP outlines a 20-year plan to direct sustainable groundwater
36 management activities that considers the needs of all users in the Basin and ensures a
37 viable groundwater resource for beneficial use by, agricultural, residential, industrial,

38 municipal and ecological users. The initial GSP is a starting point towards achievement
39 of the sustainability goal for the Basin. Although available information and monitoring data
40 have been evaluated throughout the GSP to set sustainable management criteria and
41 define projects and management actions, there are gaps in knowledge and additional
42 monitoring requirements. Information gained in the first five years of plan implementation,
43 and through the planned monitoring network expansions, will be used to further refine the
44 strategy outlined in this draft of the GSP. The GSA will work towards implementation of
45 the GSP to meet all provisions of SGMA and will utilize available local resources, and
46 resources from State and Federal agencies to achieve this. It is anticipated that
47 coordination with other agencies that conduct monitoring and/or management activities
48 will occur throughout GSP implementation to fund and conduct this important work.
49 Additional funding required may be achieved through fees, or other means, to support
50 progress towards compliance with SGMA.

51

52 **ES-2: PLAN AREA AND BASIN SETTING (CHAPTER 2)**

53 *Chapter 2 provides an overview of the Shasta Valley Basin area. This includes*
54 *descriptions of plan area, relevant agencies and programs, groundwater conditions, water*
55 *quality, interconnected surface waters, and groundwater-dependent ecosystems. These*
56 *details inform the hydrogeologic conceptual model and water budget developed for the*
57 *Basin which will be used to frame the discussion for sustainable management criteria*
58 *(Chapter 3) and projects and management actions (Chapter 4).*

59 **Description of Plan Area (Section 2.1)**

60 ***Summary of Jurisdictional Areas and Other Features (Section 2.1.1)***

61 The Shasta Valley Basin (the Basin) is a medium priority basin located in Northern
62 California. The Basin is bounded by Mount Shasta to the South, the Klamath Mountains
63 to the west and the Cascade Range to the east and the Klamath River to the north. The
64 Basin is drained by Shasta River, a tributary to the Klamath River. The primary
65 communities in Shasta Valley are the Cities of Yreka, Weed, and Montague and the
66 census-designated places of Grenada, Carrick, Gazelle, and Edgewood. As reflected in
67 the 2012-2016 disadvantaged community (DAC) Mapping Tool, Gazelle, Granada, Weed,
68 and Yreka all qualify as severely disadvantaged communities (SDACs) and Montague
69 qualifies as a DAC based on annual median household income. Land ownership in the
70 Basin is predominantly private, with two large conservation properties, California
71 Department of Fish and Wildlife's Shasta Valley and Big Springs Ranch Wildlife Areas.
72 Agriculture is a significant land use in the Basin with pasture, alfalfa, grain and hay as the
73 primary crops.

74

75 ***Water Resources Monitoring and Management Programs (Section 2.1.2)***

76 Section 2.1.2 documents monitoring and management of surface water and groundwater
77 resources in the Basin and their relation to GSP implementation. These include federal,

78 state and local agencies and their associated activities in Shasta Valley.
79

80 ***Land Use Elements or Topic Categories of Applicable General Plans (Section 2.1.3)***

81 Applicable land use and community plans in the Basin are outlined in Section 2.1.4
82 including the County of Siskiyou General Plan, City of Weed General Plan and Yreka
83 General Plan.

84 ***Additional GSP Elements (Section 2.1.4)***

85 Well policies, groundwater use regulations and the role of land use planning agencies
86 and federal regulatory agencies in GSP implementation are outlined in Section 2.1.4.

87 **Basin Setting (Section 2.2)**

88 *Section 2.2 includes descriptions of geologic formations and structures, aquifers, and*
89 *properties of geology related to groundwater, among other related characteristics of the*
90 *Basin.*

91 ***Hydrogeologic Conceptual Model (Section 2.2.1)***


92 The hydrogeologic conceptual model encompasses the Basin setting including its
93 geographical location, climate, geology, soils, land use and water management history,
94 and hydrology (Sections 2.2.1.1 through 2.2.1.5).

95 ***Current and Historical Groundwater Conditions (Section 2.2.2)***

96 *Groundwater Elevation (2.2.2.1)*

97 Groundwater data for the Basin is entirely within the DWR CASGEM Records. The
98 majority of groundwater level data available for the Basin dates back to at least the early
99 1990s, with some data available earlier and a few with only post-2010 data. Generally,
100 groundwater level data indicated levels are stable over the full period of the record as
101 shown in a subset of five wells in **Error! Reference source not found.** Groundwater
102 levels are generally shallow in the central to west-central areas of the basin (<20-40 ft
103 below ground surface) and typically do not show seasonal or longer variations. In
104 contrast, the deeper groundwater table northwest of Gazelle shows some variation with
105 drought conditions. In the volcanic aquifers, groundwater levels have generally remained
106 stable but with increases in pumping and drought conditions (post 2019), increased
107 lowering is noted, particularly in the Pluto's Cave basalt aquifer area.

108



109

110 Figure 1: Groundwater level measurements over time in five wells, one located in each
111 hydrogeologic zone.

112

113 *Estimate of Groundwater Storage (2.2.2.2)*

114 Groundwater storage is estimated based on the model, the Shasta Watershed
115 Groundwater Model (SWGM).

116 *Groundwater Quality (Section 2.2.2.3)*

117 Based on an evaluation of Basin groundwater quality using available monitoring data (see
118 Appendix 2-C), a list of constituents of interest was generated for the Basin. This list
119 includes arsenic, benzene, boron, iron, manganese, nitrate, pH and specific conductivity.
120 Multiple known contaminated sites exist in the Basin including a leaking underground
121 storage tank (LUST) site, the Davenport Property, and three open cleanup program sites
122 in Yreka as well as six California Department of Toxic Substances Control sites.

123 *Seawater Intrusion (Section 2.2.2.4)*

124 The Basin is more than 60 miles east of the Pacific Ocean and water levels are more than
125 2,000 feet above mean seal level. Seawater intrusion is not an issue in this Basin.

126 *Land Subsidence Conditions (Section 2.2.2.5)*

127 Land subsidence is lowering of the ground surface elevation and is not known to be
128 currently or historically significant in the Basin. Subsidence in Shasta Valley, based on
129 the TRE Altamira InSAR dataset provided by DWR is within the range of -0.1 to 0.1 ft,
130 largely within the margin of error indicating the absence of significant subsidence. The
131 type of geological formations present in the basin is also suggesting that future
132 subsidence is unlikely.

133

134 *Identification of Interconnected Surface Water Systems (Section 2.2.2.6)*


135 Interconnected surface water (ISW) is defined as surface water which is connected to
136 groundwater through a continuous saturated zone. SGMA mandates an assessment of
137 the location, timing, and magnitude of ISW depletions, and to demonstrate that projected
138 ISW depletions will not lead to significant and undesirable results for beneficial uses and
139 users of groundwater.

140 The Shasta River and its major tributaries are all considered part of the interconnected
141 surface water system in the Basin (Figure 2). Their large seasonal flow variations exhibit
142 all five elements of the recently proposed functional flows framework for managing
143 California rivers: fall flush flow, winter storm flow, winter baseflow, spring recess, and
144 summer baseflow. The system is also subject to significant interannual variations in flow
145 and largely affected by the complex springs system that is present throughout the valley
146 as a result of the volcanic origin.

147 The magnitude and direction of flow exchanged between surface water and groundwater
148 varies both in time and spatially (i.e., the geographic distribution of gaining and losing
149 stream reaches is not constant). When this flux is net positive into the aquifer over the
150 Basin, it is commonly referred to as stream leakage; when it is net positive into the stream
151 it is referred to as groundwater discharge.

152 In most years, the net direction in the entire watershed of stream-aquifer flux is as aquifer
153 recharge into the river, with the largest net groundwater replenishment from streams
154 occurs in wet years. Seasonally, the magnitude of leakage from the streamflow system
155 to the aquifer is greatest during late winter and early spring, while the net magnitude of
156 groundwater discharge to the stream is greatest in late fall at the end of the dry season
157 (least seasonal recharge). Spatially, the mainstem Shasta River is alternately gaining and
158 losing depending on the season, on the location, and on the year type. In other words,
159 river water weaves in and out of the aquifer on its journey along the valley floor. The upper
160 sections of tributaries tend to be losing stream reaches but conditions depend on
161 precipitation levels during any given water year and some of the tributaries tends to be
162 dry in the summer months before connecting to the main stem of the Shasta river.

163 With respect to the functional flows of the Shasta River, depletion of surface water due to
164 groundwater pumping affects the timing of the late spring recess, the amount of summer
165 baseflow, and the onset of fall flush flow.



166
167 Figure 2: Major interconnected surface waters in Shasta Valley
169

170 *Identification of Groundwater Depended Ecosystems (Section 2.2.1.6)*


171 SGMA refers to GDEs as “ecological communities or species that depend on groundwater
172 emerging from aquifers or on groundwater occurring near the ground surface”.

173 The habitat ranges of freshwater species in the Basin with special designations (i.e.,
174 endangered, threatened, species of special concern or on a watch list), were mapped.
175 Chinook salmon, coho salmon, steelhead trout, pacific lamprey and riparian vegetation

176 are all prioritized for management in the Basin as managing for these species addresses
177 the needs of other special-status species in the Basin. These prioritized species are
178 considered throughout the GSP, particularly in setting the sustainability indicators defined
179 in Chapter 3 and identifying projects and management actions identified in Chapter 4.
180 Vegetative GDE identification and classification was conducted through:

- 181 • the mapping of potential GDEs;
- 182 • assigning rooting depths based on predominant assumed vegetation type;
- 183 • establishing representations of depth to groundwater;
- 184 • identifying potential areas where depth to groundwater, rooting depth, and
185 presence of potential GDES confirm likely groundwater-dependence.

186 Potential mapped GDEs were grouped into three categories: riparian GDE, assumed
187 GDE and assumed not a GDE (where the grid-based analysis showed that the area is
188 disconnected from groundwater). Based on this analysis, around 22% of the mapped
189 potential GDE area is likely connected to groundwater and 14% of the mapped potential
190 GDE area is composed of riparian GDEs (shown in Figure 3, below).



191
192 Figure 3: Categorized GDEs for Shasta Basin.
193

194

195

196 **Water Budget (Section 2.2.3)**

197 The historical water budget for the Basin was estimated for the period October 1991
198 through 2018, using the Shasta Watershed Groundwater Model (SWGM). This 28-year
199 model period includes water years ranging from very dry (e.g., 2001 and 2014) to very
200 wet (e.g., 2006 and 2017). On an interannual scale, it includes a multi-year wet period in
201 the late 1990s and a multi-year dry period in the late 2000s and mid-2010s.

202 The water budget is presented as flows into and out of three subsystems of the integrated
203 watershed: the surface water subsystem, the soil zone (land/soil model subsystem) and
204 the groundwater subsystem.

205 Stream and lake seepage, at 124 TAF per year, accounts for 96% of the contributions
206 from surface water to the groundwater subsystem in the Basin. Fluxes from the
207 groundwater subsystem to surface water is primarily through groundwater leakage into
208 streams with an average value of 219 TAF. Agricultural pumping in the Basin accounts
209 for an average of 43 TAF per year, around one-third of the total land/soil subsystem
210 recharge in the Basin. The difference between total outflows from the groundwater
211 subsystem to land and surface water (312 TAF/ year, on average), and land and surface
212 water inflows to the groundwater subsystem (255 TAF/ year, on average) is due to net
213 groundwater inflow from outside the Basin.

214 Within the integrated model, fluxes from each subsystem to the other two subsystems are
215 simulated as distinct components (e.g. stream leakage, recharge through the soil zone,
216 and applied irrigation water). This section contains a description of each water budget
217 component.

218 Fifty-year future projected water budgets were developed using historical hydroclimate
219 data (for water years 1991-2011) and four climate change scenarios were applied to
220 explore potential effects of global warming on the Shasta Valley watershed.

221 **ES-3: SUSTAINABLE MANAGEMENT CRITERIA (CHAPTER 3)**

222 *Chapter 3 builds on the information presented in the previous Chapters and details the*
223 *key sustainability criteria developed for the GSP and associated monitoring networks.*

224 **Sustainability Goal and Sustainability Indicators (Section 3.1)**

225 **The Sustainability Goal of the Basin is to maintain groundwater resources in ways**
226 **that best support the continued and long-term health of the people, the**
227 **environment, and the economy in Shasta Valley for generations to come.**

228 The GSP details six sustainability indicators with a goal of preventing undesirable results
229 to any one of the following sustainability indicators:

230

- 231 1. Chronic Lowering of Groundwater Levels
- 232 2. Reduction of Groundwater Storage
- 233 3. Degraded Water Quality
- 234 4. Depletions of Interconnected Surface Water
- 235 5. Seawater Intrusion
- 236 6. Land Subsidence

237 Table 1 defines undesirable results for each sustainability indicator. Quantifiable minimum
 238 thresholds (MT), measurable objectives (MO), and interim milestones (IM) were also
 239 developed as checkpoints that evaluate progress made towards the sustainability goal
 240 and are quantified in Chapter 3 of the GSP. Monitoring wells throughout the basin will be
 241 used to assess conditions relevant to each sustainability indicator. Monitoring wells were
 242 selected based on well location, monitoring history, well information, and well access.

243

244 **Table 1: Shasta Valley GSP Sustainability Indicator undesirable results defined**

Sustainability Indicator	Undesirable Result Defined
Chronic Lowering of Groundwater Levels	The fall low water level observation in any of the representative monitoring sites in the Basin falls below the respective minimum threshold for 2 consecutive years.
Reduction of Groundwater Storage	Same as "Chronic Lowering of Groundwater Levels."
Degraded Water Quality	More than 25% of groundwater quality wells exceed the respective maximum threshold for concentration and/or concentrations in over 25% of groundwater quality wells increase by more than 15% per year, on average over ten years.
Depletions of Interconnected Surface Water	Greater than the depletion under which a minimum threshold of 100 CFS +/- 20% average monthly groundwater contributions occurs, for two consecutive years.
=/-Seawater Intrusion	Not applicable for the Basin.
Land Subsidence	Groundwater pumping induced subsidence is greater than the minimum threshold of 0.1 ft (0.03 m) in any single year.

245

246 **ES-4: PROJECTS AND MANAGEMENT ACTIONS TO ACHIEVE SUSTAINABILITY**
247 **(CHAPTER 4)**

248 *Chapter 4 describes past, current, and future projects management actions used to*
249 *achieve the Shasta Valley sustainability goal.*

250 To achieve the sustainability goals for Shasta Valley by 2042, and to avoid undesirable
251 results over the remainder of a 50-year planning horizon, as required by SGMA
252 regulations, multiple projects and management actions (PMAs) have been identified and
253 considered in this Groundwater Sustainability Plan (GSP).

254 Projects and management actions (PMAs) are categorized into three different tiers, as
255 follows:

256 **Tier I: Existing PMAs that are currently being implemented and are anticipated to**
257 **continue to be implemented.**

258 Projects or management actions in the Tier I category include:

- 259 • Nature Conservancy Leasing Program
- 260 • Safe Harbor Group Flow Management
- 261 • Bank Stabilization, Streambed Alteration, Floodplain Enhancement, and Riparian
262 Vegetation
- 263 • Riparian Fencing and Planting
- 264 • Novy Ice Zenkus Fish Passage Improvement Project
- 265 • Montague- Grenada Weir Modification Project
- 266 • Piezometer Transect Study Project
- 267 • City of Yreka Water Demand
- 268 • Enforcement of Survival Permits Authorizing Shasta River Template Safe Harbor
269 Agreement
- 270 • Site Plans/ Recovery of Sothern Oregon/ Northern California Coast (SONCC)
271 Coho Salmon
- 272 • Shasta River Tailwater Reduction Plan
- 273 • Upland Management

274

275 **Tier II: PMAs planned for near-term initiation and implementation (2022–2027) by**
276 **individual member agencies.**

277 Tier II PMAs include:

- 278 • Aquifer Characterization Analysis
- 279 • Avoiding Significant Increase of Total Net Groundwater Use from the basin
- 280 • Upslope Water Yield Projects
- 281 • Habitat Improvement in Shasta Watershed
- 282 • Instream Flow Leases

- 283 • Irrigation Efficiency Improvements
- 284 • Juniper Removal
- 285 • Reporting of Pump Volumes
- 286 • Voluntary Managed Land Repurposing
- 287 • Shasta Recharge Pilot Project
- 288

289 **Tier III: Additional PMAs that may be implemented in the future, as necessary**
290 **(initiation and/or implementation 2027–2042).**

291 Tier III PMAs, identified as potential future options, include:

- 292 • Alternative, lower ET Crops
- 293 • MAR and ILR
- 294 • Strategic Groundwater Pumping Curtailment
- 295 • Reservoirs

296 Additionally, other management actions are outlined that may be explored during GSP
297 implementation are outlined.

298

299 **ES-5: PLAN IMPLEMENTATION, BUDGET AND SCHEDULE (CHAPTER 5)**

300 *Section 5 details key GSP implementation steps and timelines. Cost estimates and elements of*
301 *a plan for funding GSP implementation are also presented in this section.*

302 Implementation of the GSP will focus on the following several key elements:

- 303 1. GSA management, administration, legal and day-to-day operations.
- 304 2. Implementation of the GSP monitoring program activities.
- 305 3. Technical support, including SVIHM model updates, SMC tracking, and other
306 technical analysis.
- 307 4. Reporting, including preparation of annual reports and 5-year evaluations and
308 updates.
- 309 5. Implementation of PMAs
- 310 6. Ongoing outreach activities to stakeholders

311 Total estimated annual costs for Shasta Valley Basin are between \$ 168,750 and
312 \$287,500 per year. Identified funding mechanisms include XX. The GSA may pursue
313 funding from state and federal sources for GSP implementation. As the GSP
314 implementation proceeds, the GSA will further evaluate funding mechanisms and fee
315 criteria and may perform a cost-benefit analysis of fee collection to support consideration
316 of potential refinements.

317

318

319

320

321

322